

**IN VITRO EFFECT OF ESSENTIAL OILS FROM AROMATIC AND MEDICINAL PLANTS
ON MUSHROOM PATHOGENS: *VERTICILLIUM FUNGICOLA* VAR. *FUNGICOLA*,
MYCOGONE PERNICIOSA, AND *CLADOBOTRYUM* SP.**

BRANKICA TANOVIĆ¹, IVANA POTOČNIK¹, G. DELIBAŠIĆ², M. RISTIĆ³,
M. KOSTIĆ³, and MIRJANA MARKOVIĆ¹

¹*Institute of Pesticides and Environmental Protection, 11000 Belgrade, Serbia*

²*Faculty of Agriculture, University of Belgrade, 11000 Belgrade, Serbia*

³*Dr. Josif Pančić Institute for Medicinal Plant Research, 11000 Belgrade, Serbia*

Abstract — Lavender, anise, chamomile, fennel, geranium, oregano, parsley, and sage essential oils were tested for their effectiveness against mushroom pathogens: *Verticillium fungicola* var. *fungicola*, *Mycogone perniciosa*, and *Cladobotryum* sp. Isolates were exposed to the volatile phase of the oils and then ventilated in order to determine if the effect of the oil was lethal to the pathogen. Oregano and geranium oils were the most toxic, having a fungicidal effect at 0.02-0.08 µl/ml of air, depending on the pathogen. Oregano oil was characterized by high content of carvacrol and thymol, while citranelol and geraniol were the main components of geranium oil.

Key words: Essential oils, lavender, anise, chamomile, fennel, geranium, oregano, parsley, sage, antimicrobial activity

UDC 58:582.28

INTRODUCTION

Verticillium fungicola var. *fungicola* (Preuss) Hassebrauk, *Mycogone perniciosa* (Magnus) Delacroix, and *Cladobotryum* spp. (Cooke) – the causal agents of dry bubble, wet bubble, and cobweb disease – are important fungal pathogens of the button mushroom, *Agaricus bisporus* (Lange) Imbach (Grogan and Gaze, 2000; Umaret et al., 2000; Gea et al., 2003). Symptoms of dry bubble, caused by *V. fungicola* var. *fungicola*, vary depending on the time of infection. Infection at an early stage in mushroom development results in the production of undifferentiated masses of mushrooms. If maturing mushrooms are infected, then spotting symptoms develop (Grogan et al., 2000; Potocnik et al., 2008). *Agaricus bisporus* fruit bodies infected with *M. perniciosa* become large and irregular, and tumorous fungal masses are formed (Umar et al., 2000). Exudation of accumulated extracellular fluid is observed on the surface of diseased mushrooms (Stanunton and Dunne, 1990). Caused by *Cladobotryum* spp.,

cobweb disease is characterized by growth of coarse mycelium covering affected mushrooms. As it ages, the white mycelium becomes pink (Fletcher et al., 1989).

Control of mycopathogens is based on the use of chemicals, cultural practices, and sanitation. The pesticides most commonly used on mushroom farms in Serbia are: benomyl, zinc-ethyl-enbisdithiocarbamate, and prochloraz-manganese. According to Fletcher and Yarham (1976), *V. fungicola* var. *fungicola* is resistant to benomyl, a fungicide which is frequently used to control *M. perniciosa*. By the mid-1980's, the first sign of resistance of *Cladobotryum* spp. to benomyl was recorded in Ireland and Great Britain (Gaze, 1995). Prochloraz-manganese is widely used in Europe (Bonnen and Hopkins, 1997), although moderately resistant isolates of *V. fungicola* var. *fungicola* have been found in Great Britain (Grogan et al., 2000) and Spain (Gea et al., 2003). Resistance to commonly used pesticides and the presence of residues of pesticides in food

impose the necessity of finding a suitable alternative. Application of substances of natural origin as crop protectants could be a convenient solution, safe for both human health and the environment.

Antimicrobial properties of certain essential oils have already been known for a long time (Chamberlain, 1887), but their efficacy against mycopathogenic fungi has not been well documented. Essential oils isolated from savory (*Satureja thymbra*) and sage (*Salvia pomifera* ssp. *calycina*) were investigated for antifungal activity against *M. perniciosa*; the oil of *S. thymbra* expressed better antifungal activity against *M. perniciosa* than *S. pomifera* oil (Glamoclija et al., 2006). Previous *in vitro* experiments (Tanovic et al., 2007) showed that the volatile phase of certain essential oils such as those of Scots pine, eucalyptus, juniper, orange, rosemary, and thyme, applied at a concentration of 0.65 µl/ml of air, inhibited mycelial growth of soil-borne pathogens: *Fusarium* spp., *Rhizoctonia* sp., and *Pythium* sp. In addition, application of thyme essential oil at a concentration of 1000 µl/l effectively controlled cucumber dumping-off disease (Tanovic et al., 2004) and dry bubble disease of button mushrooms caused by *V. fungicola* var. *fungicola* (Potocnik et al., 2005).

Accordingly, the objectives of this study were to investigate antimicrobial activity of several essential oils against *V. fungicola* var. *fungicola*, *M. perniciosa*, and *Cladobotryum* sp. *in vitro* and determine chemical composition of the oils expressing the highest antimicrobial activity.

MATERIALS AND METHODS

Test organisms

Isolates of *V. fungicola* var. *fungicola*, *M. perniciosa*, and *Cladobotryum* sp. identified during a 2002-2003 survey on mushroom farms in Serbia were chosen for this study.

Test substances

Essential oils of plants including lavender (*Lavandula officinalis*), anise (*Pimpinella anisum*), fennel (*Foeniculum vulgare*), geranium (*Pelargonium*

graveolens), oregano (*Origanum heracleoticum*), parsley (*Petroselinum aetheroideum*), and sage (*Salvia officinalis*) were provided by the Dr. Josif Pančić Institute for Medicinal Plant Research in Belgrade, Serbia.

Inoculum preparation

The fungal pathogens *V. fungicola* var. *fungicola*, *M. perniciosa*, and *Cladobotryum* sp. were prepared as a conidial suspension (approximately 10⁶ conidia/ml). The isolates were initially grown for 14 days on potato-dextrose-agar (PDA) plates. Conidia were harvested by flooding the plates with 10 ml of sterile distilled water and Tween 20 (v/v 0.01%), followed by filtration through a double layer of cheesecloth.

Toxicity of essential oils 'in vitro'

Antifungal activity was tested on PDA in glass Petri plates (R = 90 mm) inoculated with the investigated strains by pipetting 20 µl of the conidial suspension into a well cut in the center of the plate (R = 10 mm). The isolate was exposed to the volatile phase of essential oils for seven days at 20°C. The oils were applied as a drop onto the inner side of plate covers at concentrations of 0.02, 0.04, 0.08, 0.16, and 0.32 µl/ml of the air inside the Petri plates using a micropipette. The bottoms of the plates were immediately placed on the covers. The plates were left up-side-down and sealed by parafilm to prevent gas exchange with the outside environment. Inhibition of the mycelial growth was estimated four days after the treatment by measuring radial growth of the isolate treated with different concentrations of the oils and compared to the control. Seven days after the treatment, the plates were observed for initial mycelial growth without measuring. Concentrations of an oil which completely inhibited mycelial growth after seven-day exposure at 20°C were considered to be fungistatic and the lowest of these concentrations was determined as the minimum inhibitory concentration (MIC). Afterwards, the plates were opened and ventilated in a laminar flow hood for 30 min in order to remove volatiles and determine the fungicidal effect. The concentrations of oil were considered to be fungicidal if microbial growth was not observed seven days after ventilation. The lowest concentration with fungicidal effect was defined

as the minimum fungicidal concentration (MFC). Four replicates per treatment were used and the experiment was repeated twice.

Analysis conditions

Qualitative and quantitative analysis of essential oils was performed by gas chromatography using two detector types. A Hewlett-Packard gas chromatograph (HP-5890 Series II) was equipped with a split-splitless injector, an HP-5 capillary column (25 m, 0.32 mm i.d., 0.52 μm film thickness), and a flame ionization detector (FID). Injector and detector temperatures were set to 250 and 300°C, respectively, while the hydrogen flow rate was 1 ml/min⁻¹. Column temperature was programmed linearly in the 40-260°C temperature range at 4°C/min. Analyses with a mass spectrometer (MS) as detecting device were conducted using an HP G 1800C Series II analytical system. The same temperature program was used, while separation was performed by an HP-5MS column (30 m, 0.25 mm i.d., 0.25 μm). Helium was utilized as a carrier gas and the MS transfer line temperature was set to 260°C. The mass

detector was operated in the electron impact (EI) mode (70 eV; 40-400 m/z range). Ethanol solutions (1%) of oil samples (1 μl) were injected in a split mode (split ratio of 1: 60). Identification of essential oil components was performed using various mass spectral libraries (NIST/Wiley).

RESULTS

Antimicrobial activity of essential oils

The growth rate of the isolates was partially or completely inhibited by all tested essential oils applied at 0.02-0.32 $\mu\text{l/ml}$ of air. A 100% growth inhibition of these species was achieved by several oils at 0.32 $\mu\text{l/ml}$ of air after four-day exposure. The most sensitive species was *M. pernicioso*; the growth of this pathogen was fully inhibited by all the oils at 0.02 $\mu\text{l/ml}$ air after four-day exposure. Among the investigated oils, the most effective was oregano oil, which totally inhibited the growth of all three mycopathogenic isolates at 0.02 $\mu\text{l/ml}$ of air, while the same concentration of the other oils caused only partial inhibition of the pathogens. The only exception was geranium




Fig. 1. Effect of the volatile phase of essential oils on growth of *Verticillium fungicola* var. *fungicola* in vitro after four-day exposure.




Fig. 2. Effect of the volatile phase of essential oils on growth of *Cladobotryum* sp. in vitro after four-day exposure.

oil, which was lethal to *M. pernicioso* at 0.02 µl/ml of air. Inhibitory effects of the other tested oils varied depending on the pathogen (Figs. 1 and 2).

Toxicity of essential oils

The results obtained seven days after oil applica-

tion confirmed that the oils varied in their toxicity to *V. fungicola* var. *fungicola*, *M. pernicioso*, and *Cladobotryum* sp. isolates (Table 1). The obtained MIC and MFC values of the investigated oils ranged from 0.02 to more than 0.32 µl/ml of air. Oregano essential oil exhibited the highest level of toxicity to

Table 1. Toxicity of essential oils to *Verticillium fungicola* var. *fungicola*, *Mycogone pernicioso*, and *Cladobotryum* sp. ¹The minimal concentration of the essential oil causing complete inhibition of mycelial growth after seven-day exposure (minimum inhibitory concentration). ²The minimal concentration of the oil showing a lethal effect on the pathogen (minimum fungicidal concentration).

Essential oils	Effective concentrations of essential oils (µl/ml of air)					
	<i>Verticillium fungicola</i> var. <i>fungicola</i>		<i>Mycogone pernicioso</i>		<i>Cladobotryum</i> sp.	
	MIC ¹	MFC ²	MIC ¹	MFC ²	MIC ¹	MFC ²
Oregano (<i>Origanum heracleoticum</i>)	0.02	0.02	0.02	0.02	0.02	0.02
Geranium (<i>Pelargonium graveolens</i>)	0.08	0.16	0.02	0.02	0.04	0.08
Fenchel (<i>Foeniculum vulgare</i>)	0.08	0.32	0.04	0.04	0.16	0.32
Lavender (<i>Lavandula officinalis</i>)	0.32	0.32	0.08	0.32	0.32	> 0.32
Anise (<i>Pimpinella anisum</i>)	0.16	> 0.32	0.04	0.04	0.08	> 0.32
Parsley (<i>Petroselinum aetheroleum</i>)	> 0.32	> 0.32	0.02	0.04	0.32	0.32
Sage (<i>Salvia officinalis</i>)	> 0.32	> 0.32	0.08	0.16	> 0.32	> 0.32
Chamomile (<i>Chamomilla aetheroleum</i>)	> 0.32	> 0.32	0.16	0.16	> 0.32	> 0.32

Table 2. Chemical composition of oregano essential oil.

Component	Composition (%)
Carvacrol	65.0
Thymol	14.8
β -Phellandrene	4.3
<i>p</i> -Cymene	1.9
α -Pinene	1.4
β -Caryophyllene	1.4
α -Terpinene	1.2
γ -Terpinene	0.6
1-Octen-3-ol	0.6
β -Myrcene	0.6
Linalool	0.6
α -Thujone	0.6
Terpinen-4-ol	0.6
β -bisabolen	0.6
Borneol	0.4
Total	94.6

all isolates tested, with an MFC value of 0.02 μ l/ml, following by geranium oil, whose MFC value ranged from 0.02 to 0.16 μ l/ml of air. Lavender, chamomile, and sage oils, applied at 0.16 μ l/ml of air, did not exert a lethal effect on any of the isolates (Table 1).

Essential oil composition

Among 45 components detected in oregano essential oil, 15 constitute almost 95% of the oil mass (Table 2), while concentrations of the others are in the 0.06-0.7% range. Carvacrol and thymol were the dominant components, exceeding 79% of the oil.

In geranium oil, 54 components were identified, 27 of which were present in low concentration (less than 0.3%). Table 3 presents only components which participated in the mixture in percentages higher than 0.4%, such components constituting 91.7% of the oil.

DISCUSSION

Our results indicated that some essential oils had an ability to suppress growth of *V. fungicola* var. *fungicola*, *M. perniciosa*, and *Cladobotryum* sp. isolates *in*

Table 3. Chemical composition of geranium essential oil.

Component	Composition (%)
Citronelol	24.4
Geraniol	14.3
Linalool	12.5
Citronellyl formiate	8.5
Menthone	6.2
Geranyl-formiate	4.4
3,7-gvaiedien	4.0
α -Terpineol	3.5
Izomenthon	3.4
β -Burbonen	1.1
Tetrahydrogeraniol	1.1
α -Pinene	1.0
Geranyl-butyrate	0.8
Linalyl-propionate	0.8
Cis-rozokside	0.8
Geranyl-tiglate	0.7
β -Caryophyllene	0.7
Citronellyl-propionate	0.6
Citronellyl-butyrate	0.6
Calamenen	0.6
Neryl-propionate	0.5
Benzylidene camphor	0.4
Geranyl-propionate	0.4
δ -Gvaian	0.4
Total	91.7

vitro. Among the eight essential oils analyzed, those of oregano and geranium expressed the strongest antifungal activity against all of the investigated mycopathogens. Essential oils had previously been reported to have antimicrobial effects (Wilson et al., 1997; Suhr and Nielsen, 2003; Tanovic et al., 2004; Knezevic-Vukcevic et al., 2005; Mitic-Culafic et al., 2005; Ciric et al., 2008; Dzamic et al., 2008; Stanojevic et al., 2008). For instance, Daferera et al. (2003) reported strong activity of essential oils against the phytopathogenic fungi *Botrytis cinerea* and *Fusarium* sp. A fungistatic effect of certain oils on *Fusarium culmorum* and *Alternaria alternata* has also been demonstrated (Byron and Hall, 2002). Studies of the antifungal activity of several essential oils (Tanovic et al., 2004, 2007) against soil-borne plant pathogens including *Pythium* sp., *Verticillium albo-atrum*, and *Rhizoctonia* sp. showed that some of them had a strong inhibitory effect. Wilson et al. (1997) recorded that among the 49 essential oils tested, red thyme, cinnamon leaf, and clove bud oils showed the highest antifungal activity against *B. cinerea*, followed by oregano oil with a satisfactory effect. Oregano oil was also effective in suppress-

ing fumonisin B₁ production by *F. proliferatum* in maize grain (Velluti et al., 2003). Furthermore, this oil was highly effective in controlling internal wheat fungi in *in vivo* experiments (Paster et al., 1995). Screening experiments with 11 essential oils including those of oregano and sage against *Bacillus cereus*, a pathogen associated with food-borne disease of humans caused by toxins, showed oregano oil to be an effective growth inhibitor (Valero and Salmeron, 2003). These results demonstrated a broad spectrum of activity of oregano oil in suppressing microbial growth and indicated possible use of the oil in integrated management of various diseases.

Natural plant-derived fungicides should provide a wide variety of compounds as alternatives to synthetic fungicides, ones safe for both human health and the environment (Cutler and Hill, 1994; Daferera et al., 2003). Although the cost-effectiveness of oil application must be taken into account, this study justifies further research on practical use of essential oils for the control of *V. fungicola* var. *fungicola*, *M. perniciosus*, and *Cladobotryum* sp.

Acknowledgment — This study was carried out as part of project TR 20036, which was financially supported by the Ministry of Science and Technological Development of the Republic of Serbia.

REFERENCES

- Bonnen, A. M., and C. Hopkins (1997). Fungicide resistance and population variation in *Verticillium fungicola*, a pathogen of the button mushroom, *Agaricus bisporus*. *Mycol. Res.* **101**, 89-96.
- Byron, E. M., and A. M. Hall (2002). Inhibition of common cereal pathogenic fungi by clove oil and eucalyptus oil. *Proceedings of BCPC: Pests and Diseases*, 765-768. Farnham, Surrey.
- Chamberlain, M. (1887). Les essences au point de vue de leurs propriétés antiseptiques. *Ann. Inst. Pasteur* **1**, 153-164.
- Ciric, A., Vinterhalter, B., Savikin-Fodulovic, K., Sokovic, M., and D. Vinterhalter (2008). Chemical analysis and antimicrobial activity of methanol extracts of celandine (*Chelidonium majus* L.) plants growing in nature and cultured *in vitro*. *Arch. Biol. Sci. (Belgrade)*, **60**, 7P.
- Cutler, H. G., and R. A. Hill (1994). Natural fungicides and their delivery systems as alternatives to synthetics, In: *Biological Control of Post-harvest Diseases – Theory and Practice* (Eds. C. L. Wilson and M. E. Wisniewski), CRC Press, Boca Raton.
- Daferera, D. J., Ziogas, B. N., and M. G. Polissiou (2003). The effectiveness of plant essential oils on the growth of *Botrytis cinerea*, *Fusarium* sp., and *Clavibacter michiganensis* subsp. *michiganensis*. *Crop Protect.* **22**, 39-44.
- Dzamic, A., Sokovic, M., Ristic, M., Grujic-Jovanovic, S., Vukojevic, J., and P. D. Marin (2008). Chemical composition and antifungal activity of *Salvia sclarea* (Lamiaceae) essential oil. *Arch. Biol. Sci. (Belgrade)* **60**, 233-239.
- Fletcher, J. T., White, P. F., and R. H. Gaze (1989). *Mushrooms: Pest and Disease Control*, 58-63. Intercept, Andover, Hants.
- Fletcher, J. T., and D. J. Yarham (1976). The incidence of benomyl tolerance in *Verticillium fungicola*, *Mycogone perniciosus*, and *Hypomyces rosellus* in mushroom crops. *Ann. Appl. Biol.* **84**, 343-353.
- Gaze, R. H. (1995). *Dactylium* or cobweb. *Mushroom J.* **546**, 23-24.
- Gea, F. J., Tello, J. C., and M. J. Navarro (2003). Occurrence of *Verticillium fungicola* var. *fungicola* on *Agaricus bitorquis* mushroom crops in Spain. *J. Phytopathol.* **151**, 98-100.
- Glamoclija, J., Sokovic, M., Vukojevic, J., Milenkovic, I., and L. J. L. D. Van Griensven (2006). Chemical composition and antifungal activities of essential oils of *Satureja thymbra* L. and *Salvia pomifera* ssp. *calycina* (Sm.) Hayek. *J. Essent. Oil Res.* **18**, 115-117.
- Grogan, H. M., and R. H. Gaze (2000). Fungicide resistance among *Cladobotryum* spp. – causal agent of cobweb disease of the edible mushroom *Agaricus bisporus*. *Mycol. Res.* **104**, 357-364.
- Grogan, H. M., Keeling, C., and A. A. Jukes (2000). *In vivo* response of the mushroom pathogen *Verticillium fungicola* (dry bubble) to prochloraz-manganese, In: *BCPC Conference – Pests and Disease*, 273-278. Brighton.
- Knezevic-Vukcevic, J., Vukovic-Gacic, B., Stevic, T., Stanojevic, J., Nikolic, B., and D. Simic (2005). Antimutagenic effect of essential oil of sage (*Salvia officinalis* L.) and its fractions against UV-induced mutations in bacterial and yeast cells. *Arch. Biol. Sci. (Belgrade)* **57**, 163-172.
- Mitic-Culafic, D., Vukovic-Gacic, B., Knezevic-Vukcevic, J., Stankovic, S., and D. Simic (2005). Comparative study of the antibacterial activity of volatiles from sage (*Salvia officinalis* L.). *Arch. Biol. Sci. (Belgrade)* **57**, 173-180.
- Paster, N., Menasherov, M., Ravid, U., and B. Juven (1995). Antifungal activity of oregano and thyme essential oils applied as fumigants against fungi attacking stored grain. *J. Food Protect.* **58**, 81-85.
- Potocnik, I., Tanovic, B., Miliasevic, S., Rekanovic, E., and B. Todorovic (2005). Response of the mushroom pathogen *Verticillium fungicola* (Preuss) Hasebrauk (dry bubble) to some essential oils. *Rev. Cytol. Biol. Végét. Bot.* **28** (s.i.), 388-392.
- Potocnik, I., Vukojevic, J., Stajic, M., Tanovic, B., and B. Todorovic (2008). Fungicide sensitivity of selected *Verticillium fun-*

- gicola* isolates from *Agaricus bisporus* farms. *Arch. Biol. Sci. (Belgrade)* **60**, 151-157.
- Stanojevic, J., Beric, T., Opacic, B., Vukovic-Gacic, B., Simic, D., and J. Knezevic-Vukcevic (2008). The effect of essential oil of basil (*Ocimum basilicum* L.) on UV-induced mutagenesis in *Escherichia coli* and *Saccharomyces cerevisiae*. *Arch. Biol. Sci. (Belgrade)* **60**, 93-103.
- Staunton, L., and R. Dunne (1990). *Diseases, Molds, Disorders, and Pests of Mushrooms*. TEAGASC, Agriculture and Food Development Authority, Kinsealy Research Center, Dublin.
- Suhr, K. I., and P. V. Nielsen (2003). Antifungal activity of essential oils evaluated by two different application techniques against rye bread spoilage fungi. *J. Appl. Microbiol.* **94**, 665-674.
- Tanovic, B., Milijasevic, S., and A. Obradovic (2007). *In vitro* effect of plant essential oils on growth of some soil-borne pathogens. *Acta Hort.* **729**, 483-487.
- Tanovic, B., Obradovic, A., and I. Potocnik (2004). Effects of thyme essential oil on *Pythium* sp., In: *Proceedings of ESNA XXXIV Annual Meeting*, 502-505. Novi Sad.
- Umar, M. H., Geels, F. P., and L. J. L. D. Van Griensven (2000). Pathology and pathogenesis of *Mycogone perniciosa* infection of *Agaricus bisporus*, In: *Proceedings of the 15th International Congress on the Science and Cultivation of Edible Fungi, Maastricht, Netherlands*, 15-19 May, **2**, 561-567.
- Valero, M., and M. C. Salmeron (2003). Antibacterial activity of 11 essential oils against *Bacillus cereus* in tyndalized carrot broth. *Int. J. Food Microbiol.* **85**, 73-81.
- Velluti, A., Sanchis, V., Ramos, A. J., Egado, J., and S. Marin (2003). Inhibitory effect of cinnamon, clove, lemongrass, oregano, and palmarose essential oils on growth and fumonisin B1 production by *Fusarium proliferatum* in maize grain. *Int. J. Food Microbiol.* **89**, 145-154.
- Wilson, C. L., Soalr, J. M., El Ghaouth, A., and M. E. Wisniewski (1997). Rapid evaluation of plant extracts and essential oils for antifungal activity against *Botrytis cinerea*. *Plant Dis.* **81**, 204-210.

ДЕЛОВАЊЕ ЕТАРСКИХ УЉА ИЗ АРОМАТИЧНИХ И ЛЕКОВИТИХ БИЉАКА НА МИКОПАТОГЕНЕ ГЉИВЕ: *VERTICILLIUM FUNGICOLA* VAR. *FUNGICOLA*, *MYCOGONE PERNICIOSA* И *CLADOBOTRYUM SP. IN VITRO*

БРАНКИЦА ТАНОВИЋ¹, ИВАНА ПОТОЧНИК¹, Г. ДЕЛИБАШИЋ², М. РИСТИЋ³,
М. КОСТИЋ³, и МИРЈАНА МАРКОВИЋ¹

¹Институт за пестициде и заштиту животне средине, 11000 Београд, Србија

²Пољопривредни факултет Универзитета у Београду, 11000 Београд, Србија

³Институт за проучавање лековитог биља „Др Јосиф Панчић“, 11000 Београд, Србија

У раду су приказани резултати *in vitro* тестирања ефеката етарских уља: лаванде, аниса, камилице, морача, геранијума, оригана, першуна и жалфије, на микопатогене гљиве: *Verticillium fungicola* var. *fungicola*, *Mycogone perniciosa* и *Cladobotryum sp.* Изолати су излагани дејству пара проучаваних уља, а затим проветравани, након чега је оцењиван ефекат њихо-

вог деловања. Уља оригана и геранијума испољила су највећу токсичност за све проучаване патогене, са вредностима минималне фунгицидне концентрације између 0.02-0.08 µl/ml ваздуха, зависно од патогена. Уље оригана имало је висок садржај карвакола и тимола, док су цитранелол и гераниол биле доминантне компоненте уља геранијума.